

Students must remember the following equations:

In solving quantitative problems, students should be able to recall and apply the following equations, using standard SI units.

Equations required for Higher Tier papers only are indicated by HT in the left-hand column.

Equation number	Word equation	Symbol equation
1	weight = mass × gravitational field strength ( $g$ )	$W = m g$
2	work done = force × distance (along the line of action of the force)	$W = F s$
3	force applied to a spring = spring constant × extension	$F = k e$
4	distance travelled = speed × time	$s = v t$
5	acceleration = $\frac{\text{change in velocity}}{\text{time taken}}$	$a = \frac{\Delta v}{t}$
6	resultant force = mass × acceleration	$F = m a$
7 HT	momentum = mass × velocity	$p = m v$
8	kinetic energy = $0.5 \times \text{mass} \times (\text{speed})^2$	$E_k = \frac{1}{2} m v^2$
9	gravitational potential energy = mass × gravitational field strength ( $g$ ) × height	$E_p = m g h$
10	power = $\frac{\text{energy transferred}}{\text{time}}$	$P = \frac{E}{t}$
11	power = $\frac{\text{work done}}{\text{time}}$	$P = \frac{W}{t}$
12	efficiency = $\frac{\text{useful output energy transfer}}{\text{total input energy transfer}}$	
13	efficiency = $\frac{\text{useful power output}}{\text{total power input}}$	
14	wave speed = frequency × wavelength	$v = f \lambda$
15	charge flow = current × time	$Q = I t$
16	potential difference = current × resistance	$V = I R$
17	power = potential difference × current	$P = V I$
18	power = (current) <sup>2</sup> × resistance	$P = I^2 R$
19	energy transferred = power × time	$E = P t$
20	energy transferred = charge flow × potential difference	$E = Q V$
21	density = $\frac{\text{mass}}{\text{volume}}$	$\rho = \frac{m}{V}$

The following are given to students in the examination

Students should be able to select and apply the following equations from the *Physics equation sheet*.

Equations required for higher tier papers only are indicated by HT in the left-hand column.

Equation number	Word equation	Symbol equation
1	(final velocity) <sup>2</sup> – (initial velocity) <sup>2</sup> = 2 × acceleration × distance	$v^2 - u^2 = 2 a s$
2	elastic potential energy = 0.5 × spring constant × (extension) <sup>2</sup>	$E_e = \frac{1}{2} k e^2$
3	change in thermal energy = mass × specific heat capacity × temperature change	$\Delta E = m c \Delta \theta$
4	period = $\frac{1}{\text{frequency}}$	
5 HT	force on a conductor (at right angles to a magnetic field) carrying a current = magnetic flux density × current × length	$F = B I l$
6	thermal energy for a change of state = mass × specific latent heat	$E = m L$
7 HT	potential difference across primary coil × current in primary coil = potential difference across secondary coil × current in secondary coil	$V_s I_s = V_p I_p$